

The Hydrocarbon Potential of the Birdbear Formation, South Eastern Saskatchewan – Preliminary Results

Kathryn M. Fiess, Saskatchewan Ministry of Energy and Resources, Regina, Saskatchewan
kathryn.fiess@gov.sk.ca

GeoConvention 2012: Vision

The Upper Devonian age Birdbear Formation of south eastern Saskatchewan is comprised of porous and permeable dolostone, dolomitic limestone and anhydrite and is the stratigraphic equivalent to the Nisku Formation of Alberta (Figure 1). Light oil with gravities that range from 34 to 39 deg API has been found structurally trapped in dolostone reservoirs at Hummingbird, Kisbey, Ceylon, Flat Lake and Tatagwa (Figure 2). No stratigraphic traps have been identified for this formation in Saskatchewan to date although it is likely that such traps do exist.

The frequent inability to identify the presence of hydrocarbons from the analysis of geophysical well logs and the nebulous nature of DST results over the Birdbear carbonate, coupled with the small recoverable reserves estimates and high initial percentage water cuts for existing pools has rendered the Birdbear a less than glamorous exploration target. The geophysical well log characteristics of wells currently producing oil from Birdbear carbonates are anomalous. Oil is being produced from intervals that appear “wet” on resistivity logs (Figure 3). Furthermore, wells that recovered salt water and oil cut mud from drill stem tests are capable of significant oil production (Figure 3). The purpose of Saskatchewan Energy and Resources Birdbear Formation study of south eastern Saskatchewan is to evaluate the undeveloped hydrocarbon potential for this interval in the area defined by TWP 001 to 22 RNG 30W1M to RNG 1W3M (Figure 2). A significant aspect of the study will be to develop an understanding of the cause(s) of the low resistivity log response in the oil bearing carbonate and the possible cause(s) for salt water recoveries from drill stem test of intervals that are later perforated and capable of oil production. These data will then be assessed to determine if areas of bypassed pay potential can be identified and mapped for the Birdbear within the study area. A secondary objective is to identify area(s) where stratigraphic traps are most likely to occur.

Analytical methods to be employed in this study include log evaluations of the Birdbear Formation, water saturation calculations, analysis of production characteristics of Birdbear oil pools, core and thin section evaluation of perforated intervals with Birdbear hydrocarbon production, SEM studies and regional geological mapping. Work was initially commenced on this project in November of 2011 and the project estimated completion date is approximately November 2012. The purpose of this paper is to present a snap shot of the results to date. Thus far, the data supports the hypothesis that the Birdbear carbonate of south-eastern Saskatchewan predominantly produces oil from oil-water transition zones located on structurally closed highs. The results of preliminary thin section evaluations also support the interpretation that most of the oil and water producing Birdbear carbonate is characterized by a significant volume of microporosity. The presence of microporosity in the reservoir could substantially

reduce deep resistivity well log readings and cause anomalously high water saturation calculations for hydrocarbon bearing intervals.

Acknowledgments

Thanks are extended to Melinda Yurkowski of Saskatchewan Energy and Resources for providing me with the opportunity to work on this project and Fran Haidl for editing suggestions for this abstract.




Figure 1: Devonian Stratigraphy A) Stratigraphic chart SE Saskatchewan. B) E to W stratigraphic section of the Saskatchewan Group - Saskatchewan to Alberta.




Figure 2. Distribution of Birdbear Oil Pools in S.E. Saskatchewan.




Figure 3. Resistivity (Laterolog) and porosity (CNLDT) logs for Flat Lk. 13-03-001-16W2 well with low resistivity oil production. No free oil was recovered from the drill stem test.