


Introduction

The Winnipeg Formation is a Late Ordovician siliciclastic unit that occurs subsurface SE Saskatchewan. The formation was deposited in the Williston Basin which covers SW Manitoba and extends farther south into the neighboring states of USA (Fig.1a). The formation is exposed along the west shore of Lake Winnipeg, but none of these outcrops show a complete stratigraphic section of this formation (Oberg, 1966). The formation represents marine sedimentation during the initial period of the northward expansion of the Late Ordovician transgression in the Williston Basin (Paterson, 1971; McCabe, 1978; Norford et al., 1994). Middle Cambrian Deadwood Formation occurs disconformably on the Precambrian basement rocks and is unconformably overlain by the Winnipeg Fm. (Fig.1b) (LeFever 1996). Upper Ordovician Yeoman Formation succeeds unconformably the Winnipeg Formation in southeast Saskatchewan. In SE Saskatchewan, the Winnipeg Formation was deposited in the northeastern edge of the intracratonic Williston Basin and consists of two members: lower sandstone-dominated Black Island Member (bioturbated quartz arenite) and upper shale-dominated Icebox member (greenish grey bioturbated shale and mudstone) (Kreis 2004). The lower member is economically important and produces a fair amount of oil in North Dakota, but less in Saskatchewan and not in Manitoba. Sedimentologic, stratigraphic, diagenetic, subsurface distribution and reservoir quality of the lower sandstone-dominated member of the formation is deemed to be essential for better understanding of the hydrocarbon potential of this unit. We present here preliminary results of an on-going research work.

Figure.1. (a) Distribution of Williston basin in the United States and Canada (Worsley and Fuzesy, 1978), and (b) Precambrian to Ordovician stratigraphy in southeast Saskatchewan province (modified from Dorador et al., 2019).



The methodology includes collection, analysis and integration of data: (1) core logging and cutting analysis, (2) wireline log analysis/ thin section (3) well summary sheets/published papers. Subsurface mapping, core stratigraphic log, and correlation are used to understand the sedimentatologic depositional style of the Winnipeg Formation within the Saskatchewan province. Study area includes over 350 well information to generate the subsurface maps (Fig. 2.a). The objective of this short paper is to understand the sedimentological attributes, subsurface distribution and the reservoir properties of the lower member of the formation.

Results and observations

Subsurface mapping:- Isopach map of upper surface of the formation and porosity distribution have been generated and interpreted. The isopach map shows that mapping is applied to investigate the member's depositional setting, thickness variations and subsurface distribution of the porous sandstone. Subsurface mapping of the total thickness of the formation shows variations in three zones (Fig.2.b). Zone 1, which constitutes most of the mapped area, shows that the formation thickens towards southeast of the province. Zone 2 occurs west of the zone 1

and shows a relatively thin distribution of the formation reaching less 8 meters. This zone shows northeast-southwest trend possibly controlled by paleotopographic feature. Farther west of zone 2, the formation get thicker westward. Depth structure map and 3D mapping (Fig.2.c & d) of the upper surface of the formation also confirms the occurrence an elevated zone which matches that of zone 2. The map also shows that the basin was shallower towards north, northeast and northwest and deep towards south, thus conforming that the depocenter of the Williston Basin was located farther south of the international border.




Figure.2. Subsurface distribution of the Winnipeg Formation (a) Study area (b) Isopach map showing SE-ward thickening of the formation. (c) 3D depth map from the upper surface of the formation. The formation deepens SE –ward. (d) 2D map of the upper surface of the formation showing subsurface distribution comparable to that of the 3D map. See text for discussion of the three zones (1, 2 & 3)

Reservoir properties:- For the reservoir characterization, porosity interval map of the Black Island Member shows that the thickness of the porous intervals is also split in three zones that match the same zones of the thickness distribution (Fig.3.a). The eastern large area, the thickness of the porous interval has an overall southeastward increase. In zone 2 area, the porous zone is the thinnest and then increases westward. The thickness distribution maps suggest existence of a paleohigh in zone 2 and, possibly pre-Yeoman Formation erosion caused the thickness reduction in zone 2.

Three wells core (131/03-08-017-19w2/00, 21/08-06-007-15w2/0, and 01/05-15-010-02w2/0) are also interpreted on the basis of facies analysis, sedimentary structures and trace fossils. Black Island Member (sandstone) mostly consist of white to light and dark grey, very fine to coarse grained containing pyrite, glauconite and quartz grains, very porous, moderate to poor

consolidation, and fair porosity are observed. Grey to brownish sandstones color is due to organic matter associated with the bioturbation and calcite cementation and early to late diagenesis is also very dominant with quartz arenite (Fig.3.b & c). Correlation is also done between three wells that indicates that Black Island Member is dominant with coarsening upward sequence indicates low sea level settings. Icebox Member is fining upward sequence with high sea level conditions. The Black Island Member thickness indicates that it is more widespread than the Icebox Member in the province




Figure.3. Porous zones estimation of Winnipeg Formation (a) Porosity interval map of Black Island Member indicates more porous zone is toward southern and western site (b) Quartz arenite with pores and interbedded grains at depth 2066.5m of 131/03-08-010-19w2/00 well, and (c) Quartz arenite with pores and interbedded grains at depth 2292.9m of 101/05-15-010-2w2/00 well.

Conclusions

The Winnipeg Formation consists of a lower quartz arenite member (Black Island Mb.) and upper shale-dominated member (Icebox Mb.) The lower member was deposited in a medium to high energy shoreface and constitutes the reservoir unit of the formation in SE Saskatchewan. 2D and 3D isopach map of the formation, structure of its upper surface map and thickness variation of the porous zone of the lower member furnish the subsurface distribution of these properties. The maps show three zones that can be summarized as follows. Zone 1: this zone occurs about 3/4th of the study area and is characterized by thick section of the formation that gets deeper SE-ward. It also preserves the thickest pay zone which also gets deeper SE-ward. Zone 2 is a relatively narrow, NE-SE-oriented corridor of apparently uplifted appearance. This

corridor constitutes the shallowest zone and thinnest porous zone. Zone 3 occurs west of zone 2 and gets thicker and deeper away from the uplifted corridor.

Acknowledgements

We thank the Saskatchewan Geological Survey for giving us access to the core lab and allowing usage of the facility in their subsurface lab. This project is also financially supported by the Ministry of Energy and Resources of Saskatchewan.

Selected references:-

- Carlson, C.G. 1960: Stratigraphy of the Winnipeg and Deadwood Formations in North Dakota. North Dakota Geological Survey Bulletin 35, 1–149
- Carlson, C.G. & Thompson, S.C. 1987: Stratigraphy of the Deadwood Formation and Winnipeg Group in the Williston Basin – anatomy of a cratonic oil province Fifth International Williston Basin Symposium, June, 1987. Grand Forks, North Dakota, USA. 71–80.
- Carlson, C.G. and Anderson, S.B., 1965, Sedimentary and tectonic history of North Dakota part of the Williston Basin: American Association of Petroleum Geologists Bulletin, v.49, No. 11, p 1833-1846.
- Dorador, J., Buatois, L.A., Mangano, M.G. & Rodriguez-Tovar, F.J. (2019) Ichnology of the Winnipeg Formation, southeast Saskatchewan: a glimpse into the marine infaunal ecology of the Great Ordovician Biodiversification Event, Paper A-4, 17 pp. Saskatchewan, Canada.
- Holmden, C. 2009: Ca isotope study of Ordovician dolomite, limestone, and anhydrite in the Williston Basin: implications for subsurface dolomitization and local Ca cycling. *Chemical Geology* 268, 180–188.
- Kendall, A.C. (1976): The Ordovician Carbonate Succession (Bighorn Group) of Southeastern Saskatchewan; Sask. Dep. Mineral Resour., Rep. 180, 185p.
- Kreis, L.K. (2004a): Geology of the Middle Cambrian–Lower Ordovician Deadwood Formation in Saskatchewan; Lower Paleozoic Map Series – Saskatchewan, Sask. Industry Resources, Misc. Rep. 2004-8, CD-ROM, Sheet 2 of 8; <http://economy.gov.sk.ca/MiscRep2004-8>.
- LeFever, R.D. (1996): Sedimentology and stratigraphy of the Deadwood-Winnipeg interval (Cambro-Ordovician)
- McCabe, H.R. (1978): Reservoir Potential of the Deadwood and Winnipeg Formations in Southwest Manitoba;
- Nesheim, T.O. (2012): Review of Ordovician Black Island Formation (Winnipeg Group) Oil and Gas Production: North Dakota Geological Survey, Geologic Investigation No. 161, 13 p
- Norford, B.S., Haidl, F.M., Bezys, R.K., Cecile, M.P., McCabe, H.R., and Paterson, D.F. (1994): Chapter 9: Middle Ordovician to Lower Devonian strata of the Western Canada Sedimentary Basin; in Mossop, G.D. and Shetsen, I. (eds), Geological Atlas of the Western Canada Sedimentary Basin; Can. Soc. Petrol. Geol./Alta. Resear. Counc., Calgary, p109-127.
- Oberg, R. 1966: Winnipeg conodonts from Manitoba. *Journal of Paleontology* 40, 130–147.
- Paterson, D.F. (1971): The Stratigraphy of the Winnipeg Formation (Ordovician) of Saskatchewan: Department of Mineral Resources, Geological Science Branch (Sedimentary Geology Division), Report No.140.,p.5.
- TGI Williston Basin Working Group (2008) Ordovician Winnipeg Formation: isopach; Manitoba Science, Technology, Energy and Mines, Manitoba Geological Survey, Stratigraphic Map SM2008-OW-I, scale 1:1 000 000.
- Worsley, N., and Fuzesy, A., 1978, The potash-bearing members of the Devonian Prairie evaporite of southeastern Saskatchewan south of the mining area in Economic Geology of the Williston Basin: Williston Basin Symposium, Montana Geol. Soc., p. 153-165.